

10/559,878

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssspta1611bxv

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	JAN 02	STN pricing information for 2008 now available
NEWS	3	JAN 16	CAS patent coverage enhanced to include exemplified prophetic substances
NEWS	4	JAN 28	USPATFULL, USPAT2, and USPATOLD enhanced with new custom IPC display formats
NEWS	5	JAN 28	MARPAT searching enhanced
NEWS	6	JAN 28	USGENE now provides USPTO sequence data within 3 days of publication
NEWS	7	JAN 28	TOXCENTER enhanced with reloaded MEDLINE segment
NEWS	8	JAN 28	MEDLINE and LMEMLINE reloaded with enhancements
NEWS	9	FEB 08	STN Express, Version 8.3, now available
NEWS	10	FEB 20	PCI now available as a replacement to DPCI
NEWS	11	FEB 25	IFIREF reloaded with enhancements
NEWS	12	FEB 25	IMSPRODUCT reloaded with enhancements
NEWS	13	FEB 29	WPINDEX/WPIDS/WPIX enhanced with ECLA and current U.S. National Patent Classification
NEWS	14	MAR 31	IFICDB, IFIPAT, and IFIUDB enhanced with new custom IPC display formats
NEWS	15	MAR 31	CAS REGISTRY enhanced with additional experimental spectra
NEWS	16	MAR 31	CA/CAPplus and CASREACT patent number format for U.S. applications updated
NEWS	17	MAR 31	LPCI now available as a replacement to LDPCI
NEWS	18	MAR 31	EMBASE, EMBAL, and LEMBASE reloaded with enhancements
NEWS	19	APR 04	STN AnaVist, Version 1, to be discontinued
NEWS	20	APR 15	WPIDS, WPINDEX, and WPIX enhanced with new predefined hit display formats
NEWS	21	APR 28	EMBASE Controlled Term thesaurus enhanced
NEWS	22	APR 28	IMSRESEARCH reloaded with enhancements
NEWS	23	MAY 30	INPAFAMDB now available on STN for patent family searching
NEWS	24	MAY 30	DGENE, PCTGEN, and USGENE enhanced with new homology sequence search option
NEWS	25	JUN 06	EPFULL enhanced with 260,000 English abstracts
NEWS	26	JUN 06	KOREAPAT updated with 41,000 documents
NEWS	27	JUN 13	USPATFULL and USPAT2 updated with 11-character patent numbers for U.S. applications
NEWS	28	JUN 19	CAS REGISTRY includes selected substances from web-based collections
NEWS	29	JUN 25	CA/CAPplus and USPAT databases updated with IPC reclassification data
NEWS	30	JUN 30	AEROSPACE enhanced with more than 1 million U.S. patent records

10/559,878

NEWS 31 JUN 30 EMBASE, EMBAL, and LEMBASE updated with additional
options to display authors and affiliated
organizations
NEWS 32 JUN 30 STN on the Web enhanced with new STN AnaVist
Assistant and BLAST plug-in
NEWS 33 JUN 30 STN AnaVist enhanced with database content from EPFULL

NEWS EXPRESS JUNE 27 08 CURRENT WINDOWS VERSION IS V8.3,
AND CURRENT DISCOVER FILE IS DATED 23 JUNE 2008.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 10:17:58 ON 25 JUL 2008

=> file caplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'CAPLUS' ENTERED AT 10:18:16 ON 25 JUL 2008
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is
held by the publishers listed in the PUBLISHER (PB) field (available
for records published or updated in Chemical Abstracts after December
26, 1996), unless otherwise indicated in the original publications.
The CA Lexicon is the copyrighted intellectual property of the
American Chemical Society and is provided to assist you in searching
databases on STN. Any dissemination, distribution, copying, or storing
of this information, without the prior written consent of CAS, is
strictly prohibited.

FILE COVERS 1907 - 25 Jul 2008 VOL 149 ISS 5
FILE LAST UPDATED: 24 Jul 2008 (20080724/ED)

Caplus now includes complete International Patent Classification (IPC)
reclassification data for the second quarter of 2008.

Effective October 17, 2005, revised CAS Information Use Policies apply.
They are available for your review at:

<http://www.cas.org/legal/infopolicy.html>

=> s (fluorination or fluoridation)

10/559,878

```
          17987 FLUORINATION
          1261 FLUORIDATION
L1        19144 (FLUORINATION OR FLUORIDATION)
```

```
=> s diphenyliodonium
L2        2148 DIPHENYLIODONIUM
```

```
=> s fluoride
L3        275699 FLUORIDE
```

```
=> s water
L4        2741080 WATER
```

```
=> s l1 adn l2 and l3 and l4
```

MISSING OPERATOR L1 ADN

The search profile that was entered contains terms or
nested terms that are not separated by a logical operator.


```
=> s l1 and l2 and l3 and l4
L5        0 L1 AND L2 AND L3 AND L4
```

```
=> s l1 and l2
L6        6 L1 AND L2
```

```
=> d l6 1-6 bib ABS
```

10/559,878

L6 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN
AN 2007:1310500 CAPLUS
DN 148:121427
TI A practical route for synthesizing a PET ligand containing
[18F]fluorobenzene using reaction of diphenyliodonium salt with
[18F]F-
AU Zhang, Ming-Rong; Kumata, Katsushi; Suzuki, Kazutoshi
CS Radiochemistry Section, Department of Molecular Probes, Molecular Imaging
Center, National Institute of Radiological Sciences, Inage-ku, Chiba,
263-8555, Japan
SO Tetrahedron Letters (2007), 48(49), 8632-8635
CODEN: TELEAY; ISSN: 0040-4039
PB Elsevier Ltd.
DT Journal
LA English
OS CASREACT 148:121427
GI



AB A practical route was developed for preparing a fluorine-18 ([18F]) labeled ligand containing a [18F]fluorobenzene ring by employing the reaction of diphenyliodonium salt with [18F]F-. Diphenyliodonium tosylate was synthesized from the appropriate tributylphenylstannane with [hydroxy(tosyloxy)iodo]benzene. Using this method, [18F]DAA1106 (I), a positron emission tomog. ligand for imaging peripheral-type benzodiazepine receptor, was prepared


RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN
 AN 2007:614337 CAPLUS
 DN 148:307964
 TI Development of molecular probe labeling technology. Practical synthesis of
 [18F]fluorobenzene with [18F] nucleophilic substitution reaction
 AU Zhang, Ming- Rong; Kumada, Katsushi; Suzuki, Kazutoshi
 CS National Institute of Radiological Sciences, Japan
 SO Hoshasen Igaku Sogo Kenkyusho, [Report] NIRS-M (2007),
 NIRS-M-197(Dai-1-kai Bunshi Imejingu Kenkyu Senta Shinpojumu, 2007), 19-24
 CODEN: NIRRDY
 PB Hoshasen Igaku Sogo Kenkyusho
 DT Journal; General Review
 LA Japanese
 AB A review on synthesis of [18F]fluorobenzene derivs., e.g. [18F]DAA1106,
 for PET ligands by generation of diphenyliodonium tosylates from
 hydroxy(4-methoxyphenyl)(4-methylbenzenesulfonato-O)iodine (CAS REG
 126550-93-4) and tributylphenyltin derivs. and regioselective nucleophilic
 substitution reaction of the resulting diphenyliodonium
 tosylates with [18F]F-.

L6 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN
AN 2006:214878 CAPLUS
DN 145:155939
TI Non-PFOS photoacid generating compounds for chemically amplified resists
AU Ayothi, Ramakrishnan; Yi, Yi; Felix, Nelson; Ober, Christopher K.; Cao, Heidi; Wang, Yueh
CS Department of Materials Science and Engineering, Cornell University, Ithaca, NY, 14853, USA
SO Polymer Preprints (American Chemical Society, Division of Polymer Chemistry) (2006), 47(1), 528-529
CODEN: ACPPAY; ISSN: 0032-3934
PB American Chemical Society, Division of Polymer Chemistry
DT Journal; (computer optical disk)
LA English
AB New class of photoacid generators (PAGs) that carry an anion composed of aryl groups and possessing less fluorination than perfluorooctane sulfonate (PFOS), thus more environmentally friendly, were developed. These include diphenyliodonium 2-(phenoxy)tetrafluoroethane-1-sulfonate (PAG 1) and diphenyliodonium 2-nitro-4-(trifluoromethyl)benzenesulfonate (PAG 2). These PAGs have good solubility in common solvents and comparable thermal properties to standard ionic PAGs. These PAGs show reasonable sensitivity upon exposure to e-beam radiation and are capable of resolving sub-100 nm lines and spaces. The two new PAGs may have a higher degradation probability but isolation of the pure acids to evaluate their mammalian toxicity, ecotoxicity and bioconcn. factor will provide a better understanding of the environmentally friendly nature of the new non-PFOS PAGs.
RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN
 AN 2005:588844 CAPLUS
 DN 143:115340
 TI Process for fluorination and radiofluorination of iodonium salts
 in the presence of a radical trap
 IN Wadsworth, Harry John; Widdowson, David Arthur; Wilson, Emmanuelle;
 Carroll, Michael Andrew
 PA GE Healthcare Limited, UK
 SO PCT Int. Appl., 32 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005061415	A1	20050707	WO 2004-GB5304	20041217
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	EP 1697279	A1	20060906	EP 2004-806112	20041217
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS				
	CN 1898184	A	20070117	CN 2004-80038469	20041217
	JP 2007515465	T	20070614	JP 2006-546303	20041217
	US 20060292060	A1	20061228	US 2006-559879	20060830
PRAI	GB 2003-29716	A	20031223		
	WO 2004-GB5304	W	20041217		
OS	CASREACT 143:115340; MARPAT 143:115340				
GI					



AB Decomposition of iodonium salts I [Q = precursor of fluorine-labeled compound; Y = anion selected from triflate, nonaflate, mesylate, hexaflate; R1-R2, R4-R5 = independently H, NO₂, CN, halo, (un)protected C1-10 hydroxyalkyl, C2-10 carboxyalkyl, C1-10 alkyl, C2-10 alkoxyalkyl, C1-10 aminoalkyl, C1-10 haloalkyl, C6-14 aryl, C3-12 heteroaryl, C3-20 alkylaryl, C5-12 arylene, C2-10 alkenyl, C2-10 alkynyl, C1-10 acyl, C7-10 aroyl, C2-10 carboalkoxy, C2-10 carbamoyl, C2-10 carbamyl, C1-10 alkylsulfinyl; or R1-R5 may form 4-6-membered ring; R3 = any group R1-R2, R4-R5 or link to a solid support] by a free radical process has been identified as a significant factor in the observed yield variability of fluoridation

reactions using said iodonium salts. Accordingly, the inclusion of a free radical trap, such as 2,2,6,6-tetramethylpiperidine-N-oxide, 1,2-diphenylethylene, ascorbate, p-aminobenzoic acid, α -tocopherol, hydroquinone, di-t-butylphenol, β -carotene, or gentisic acid in the reaction mixture blocks the radical chain decomposition pathway for iodonium salts such that only the reaction leading to fluoridation can occur and the yield of aryl fluoride becomes high and reproducible. In both the solution and the solid phase the preferred method of the present invention is radiofluoridation. Thus, radiofluorination of diphenyliodonium triflate with ^{18}F -fluoride in the presence of Kryptofix 222 in dry acetonitrile and 70 mol % 2,2,6,6-tetramethylpiperidine-N-oxide gave radiolabeled fluorobenzene in 41-57% yield and 82-96% radiochem. purity. The same reaction without the radical trap gave labeled fluorobenzene in 0-40% yields and 0-65% radiochem. purity.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN

AN 1999:96208 CAPLUS

DN 130:168015

TI Ionic perfluorosulfonimide compounds with delocalized anionic charge, and their use as components of ionic conductors or catalysts

IN Armand, Michel; Michot, Christophe; Yagupolskii, Yurii; Yagupolskii, Lev; Bezdudny, Andrej; Kondratenko, Natalya

PA Acep Inc., Can.; Universite de Montreal; Centre National de la Recherche Scientifique; Institute of Organic Chemistry

SO PCT Int. Appl., 59 pp.

CODEN: PIXXD2

DT Patent

LA French

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9905100	A1	19990204	WO 1998-FR1663	19980727
	W: CA, JP, UA, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	CA 2266643	A1	19990204	CA 1998-2266643	19980727
	EP 928287	A1	19990714	EP 1998-941464	19980727
	EP 928287	B1	20031001		
	R: DE, FR, GB, IT				
	JP 2001507043	T	20010529	JP 1999-509451	19980727
	EP 1388546	A2	20040211	EP 2003-292375	19980727
	EP 1388546	A3	20040303		
	R: DE, FR, GB, IT				
	US 6340716	B1	20020122	US 1999-269264	19990325
	US 20020013381	A1	20020131	US 2001-931076	20010817
	US 6548567	B2	20030415		
	US 20030195269	A1	20031016	US 2003-366450	20030214
	US 20040162362	A9	20040819		
	US 6841638	B2	20050111		
	US 20050158631	A1	20050721	US 2005-32038	20050111
	US 20070205388	A1	20070906	US 2007-654035	20070117
	US 7378034	B2	20080527		
PRAI	CA 1997-2211465	A	19970725		
	EP 1998-941464	A3	19980727		
	WO 1998-FR1663	W	19980727		
	US 1999-269264	A3	19990325		
	US 2001-931076	A3	20010817		
	US 2003-366450	A3	20030214		
	US 2005-32038	B1	20050111		

OS MARPAT 130:168015

AB The invention concerns ionic compds. of formula $[R_1X_1(:Z_1)Q-X_2(:Z_2)R_2]_m$ Mm^+ [I; in which Mm^+ is a cation of valence m; each $X_i = S:Z_3$, $S:Z_4$, PR_3 , or PR_4 ; $Q = N$, CR_5 , CCN , or CSO_2R_5 ; each $Z_i = :O$, $:NC.tplbond.N$, $:C(C.tplbond.N)_2$, $:NS(:Z)2R_6$, or $:C[S(=Z)2R_6]_2$; each $R_i = Y$, YO , YS , Y_2N , or F ; $Y =$ monovalent organic radical, or repeat unit of a polymeric fabric]. I are useful for preparing materials with ionic conduction, electrolytes, as catalysts for polymerization and other organic reactions, and for doping polymers.

For instance, butanesulfonyl chloride was condensed with $CF_3SO_2NH_2$ using DABCO, and the product treated with saturated KCl and AcOH to give crystalline $BuSO_2N(K)SO_2CF_3$. This was treated with $(COCl)_2$ and DMF in MeCN, followed by treatment with $CF_3SO_2NH_2$ and DABCO, and then workup with aqueous KCl and AcOH, to give title compound $CF_3SO_2N-S(:O)(Bu):NSO_2CF_3 K^+$. The latter was converted to the corresponding Li^+ salt using $LiBF_4$, and the Li salt was incorporated in poly(ethylene oxide) of mass 106 to give a film with conductivity

10/559,878

>2 + 10⁻⁵ S/cm at 25°.

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2008 ACS on STN

AN 1991:558135 CAPLUS

DN 115:158135

OREF 115:27063a,27066a

TI Direct and regiocontrolled synthesis of α -phenyl ketones from silyl enol ethers and diphenyliodonium fluoride

AU Chen, Kuanchiang; Koser, Gerald F.

CS Dep. Chem., Univ. Akron, Akron, OH, 44325, USA

SO Journal of Organic Chemistry (1991), 56(20), 5764-7
CODEN: JOCEAH; ISSN: 0022-3263

DT Journal

LA English

OS CASREACT 115:158135

AB The efficacy of diphenyliodonium fluoride (I), for the phenylation of silyl enol ethers was investigated. When the silyl enol ethers of cyclopentanone, 2-methylcyclopentanone, cyclohexanone, 2-methylcyclohexanone, acetophenone, 2-pentanone, diisopropyl ketone, and pinacolone were mixed with I in THF, either α -Ph or α,α -di-Ph ketones were produced and isolated in yields ranging from 20 to 88%. The regiochem. of α -phenylation can be controlled by appropriate choice of silyl enol ether was demonstrated. 3,3-Dimethyl-2-(silyloxy)-1-butene gave a dehydro dimer of pinacolone with I in addition to α -phenylpinacolone, thus suggesting that phenylations of silyl enol ethers with I may proceed via radical intermediates.

10/559,878

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

30.46

30.67

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-4.80

-4.80

STN INTERNATIONAL LOGOFF AT 10:21:05 ON 25 JUL 2008